

Stable and unstable perturbations in universal scaling phenomena far from equilibrium

Thimo Preis

Together with Michal P. Heller and Jürgen Berges

QSEC 2022, 17.11.2022, based on [2209.14883](#)



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386




STRUCTURES
CLUSTER OF
EXCELLENCE

Nonthermal fixed points

Universal self-similar scaling far from equilibrium:


$$f(t, \mathbf{p}) = (t/t_{\text{ref}})^\alpha f_S((t/t_{\text{ref}})^\beta \mathbf{p})$$

Berges et al., PRL 114, 061601 (2015)



See also: Kurkela et al., PRL 115, 182301 (2015),
 Micha et al., PRL 90, 121301 (2003),
 Berges et al., PRL 101, 041603 (2008).

Prüfer et al., Nature 563, 217 (2018)



See also: Erne et al., Nature 563, 225 (2018),
 Glidden et al., Nature Phys. 17, 457 (2021).

Far-from-equilibrium quantum fields

$$\hat{H}(t) = \int d^3x \left[\frac{1}{2} \left(\partial_t \hat{\Phi}_a(t, \mathbf{x}) \right)^2 + \frac{1}{2} \left(\nabla_{\mathbf{x}} \hat{\Phi}_a(t, \mathbf{x}) \right)^2 + \frac{\lambda}{4!N} \left(\hat{\Phi}_a(t, \mathbf{x}) \hat{\Phi}_a(t, \mathbf{x}) \right)^2 \right]$$

$$\langle \hat{H}(t) \rangle \longrightarrow \text{Feynman diagrams: two circles, two circles with a line, triangle, square, ...}$$


Solve quantum evolution equations of

$$F_{ab}(t, t', \mathbf{x} - \mathbf{x}') = \frac{1}{2} \langle \{ \hat{\Phi}_a(t, \mathbf{x}), \hat{\Phi}_b(t', \mathbf{x}') \} \rangle - \langle \hat{\Phi}_a(t, \mathbf{x}) \rangle \langle \hat{\Phi}_b(t', \mathbf{x}') \rangle$$

$$\rho_{ab}(t, t', \mathbf{x} - \mathbf{x}') = i \langle [\hat{\Phi}_a(t, \mathbf{x}), \hat{\Phi}_b(t', \mathbf{x}')] \rangle$$


$$F(t_0, t_0, \mathbf{p}) = \frac{1}{\sqrt{\mathbf{p}^2 + m^2}} \left[\left(\frac{N n_0}{\lambda} \right) \theta(Q - |\mathbf{p}|) + \frac{1}{2} \right]$$

We consider $\lambda = 0.01$, $N = 4$ and $n_0 = 25$ in this work.



Perturbations around the universal scaling solution

$$F(t, t, |\mathbf{p}|) = (t/t_{\text{ref}})^\alpha F_s((t/t_{\text{ref}})^\beta |\mathbf{p}|) + \delta F(t, t, |\mathbf{p}|)$$



Far-from-equilibrium quasi-particles

Wigner transform with $\tau = \frac{1}{2}(t + t')$ and $\Delta t = t - t'$

$$\rho(\tau, \omega, |\mathbf{p}|) = -i \int_{-2\tau}^{2\tau} d\Delta t e^{i\omega\Delta t} \rho(\tau + \frac{\Delta t}{2}, \tau - \frac{\Delta t}{2}, |\mathbf{p}|)$$


Tower of quasi-particles

$$\omega(t, \mathbf{p}) = \sqrt{\mathbf{p}^2 + M^2(t)}$$

$$\omega^\pm(t, \mathbf{p}) = 2M(t) \pm \omega(t, \mathbf{p})$$

Interpretation as phase fluctuations of a macroscopic zero mode rotating in N-component field space.

Boguslavski et al., PRD 101, 091902 (2020)



Linear response around the universal scaling solution

Perturbations described by time-dependent response rate:


$$\frac{\partial_t \delta F(t, \omega, |\mathbf{p}|)}{\delta F(t, \omega, |\mathbf{p}|)} = -\gamma(t, \omega, |\mathbf{p}|)$$

where

$$\gamma(t, \omega \simeq M, |\mathbf{p}| \simeq 0) < 0$$

To leading order in gradients, the diagonal contribution to the response rate is

$$\gamma(t, \omega, |\mathbf{p}|) = -\frac{\Sigma_\rho(t, \omega, |\mathbf{p}|)}{2\omega}$$



Comparison with on-shell linear response

Dynamics of perturbations are described by


$$\frac{\delta F(t, \omega, |\mathbf{p}|)}{\delta F(t_i, \omega, |\mathbf{p}|)} = e^{-\Gamma(t, \omega, |\mathbf{p}|)}$$

with rate integral

$$\Gamma(t, \omega, |\mathbf{p}|) = \int_{t_i}^t dt' \gamma(t', \omega, |\mathbf{p}|)$$

Initial kick at $t_i Q = 1800$

$$\begin{aligned} \delta F(t_i, t_i, |\mathbf{p}|) &= 10^{-3} (t_i/t_{\text{ref}})^\alpha \\ &\times F_S((t_i/t_{\text{ref}})^\beta |\mathbf{p}|) \end{aligned}$$



Interpretation via scattering processes

Elastic scatterings describe the leading contribution to the on-shell linear response rate ($p = (p^0, \mathbf{p})$)

$$\gamma^{2\leftrightarrow 2}(t, p) = -\frac{\lambda^2}{36Np^0} \int \frac{d^3q}{(2\pi)^3} \frac{d^3l}{(2\pi)^3} \frac{d^3r}{(2\pi)^3} (2\pi)^4$$

with $F(t, p) = \left[f_p(t) + \frac{1}{2} \right] \rho(t, p)$


$$\times \int_0^\infty \frac{dq^0 dl^0 dr^0}{(2\pi)^3} \delta^{(4)}(p + l - q - r) \rho(t, q) \rho(t, l) \rho(t, r)$$

$$\times [v_{\text{eff}}(t, p + l) + v_{\text{eff}}(t, p - q) + v_{\text{eff}}(t, p - r)]$$

$$\times [(f_l(t) + 1)f_q(t)f_r(t) - f_l(t)(f_q(t) + 1)(f_r(t) + 1)]$$


$v_{\text{eff}}(t, p) = \left| 1 + \frac{\lambda}{3} \int \frac{d^4k}{(2\pi)^4} F(t, p - k) G^R(t, k) \right|^{-2}$

s-channel resummation:



$$\rho(t, q) \sim \delta((q^0)^2 - (\omega(t, \mathbf{q}))^2)$$

Two-to-two scattering with $p^0 + \omega(t, \mathbf{l}) = \omega(t, \mathbf{q}) + \omega(t, \mathbf{r})$ receives contributions from



$$p^0 = \omega(t, \mathbf{p})$$

$$\gamma_+^{2\leftrightarrow 2}$$

and

$$p^0 = \omega^-(t, \mathbf{p})$$

$$= 2M(t) - \omega(t, \mathbf{p})$$

$$\gamma_-^{2\leftrightarrow 2}$$

Universal scaling of perturbations

The on-shell rate integral exhibits scaling


$$\Gamma(t, |\mathbf{p}|) = (t/t_{\text{ref}}) \Gamma_S((t/t_{\text{ref}})^\beta |\mathbf{p}|)$$

where the scaling function can be described by

$$\Gamma_S((t/t_{\text{ref}})^\beta |\mathbf{p}|) = A (t/t_{\text{ref}})^\beta |\mathbf{p}|/Q - B$$

For a given momentum $|\mathbf{p}|$, the positive (stable) contribution $\sim A t^{\beta+1} |\mathbf{p}|$ will eventually outgrow the negative (unstable) $\sim B t$ term for all non-zero momenta in the scaling regime.

Scaling instability



Summary

based on [2209.14883](#)

We studied the dynamics of perturbations around nonthermal fixed points, which revealed the presence of both stable and unstable perturbations.

Linear response theory allowed us to demonstrate that unstable dynamics arises from a competition between elastic scattering processes among different quasi-particle states.

We discovered the phenomenon of a [scaling instability](#). As a consequence, the system shows attractor behavior after $t \sim 1/|\mathbf{p}|^{1/\beta}$ for any non-vanishing momentum $|\mathbf{p}|$.

Our results demonstrate from first principles that [self-organized scaling](#) (no fine-tuning) can be realized in the presence of both stable and unstable directions for the dynamics.

Outlook

Investigate this phenomenon for non-relativistic system, access via experimental platforms with ultra-cold quantum gases in the many-body regime with sufficiently large volumes.